

Isomonodromic deformation with an irregular singularity and the Θ function

Kazuhide Matsuda

Department of Mathematics, Graduate School of Information Science and Technology,
Osaka University,
1-1 Machikaneyama-machi, Toyonaka, 560-0043.

Abstract: We will study a monodromy preserving deformation with an irregular singular point and determine the τ function of the monodromy preserving deformation by the elliptic θ function moving the argument z and the period Ω .

Key words: τ function; θ function; monodromy preserving deformation; irregular singular point.

1 Introduction

In this paper, we construct a monodromy preserving deformation whose τ function is represented by the θ function moving both the argument z and the period Ω . The monodromy preserving deformation has an irregular singularity of the Poincaré rank one.

In [3], Miwa-Jimbo-Ueno extended the work of Schlesinger in [1] and established a general theory of monodromy preserving deformation for a first order matrix system of ordinary linear differential equations,

$$\frac{dY}{dx} = A(x)Y, \quad A(x) = \sum_{\nu=1}^n \sum_{k=0}^{r_\nu} \frac{A_{\nu,-k}}{(x-a_\nu)^{k+1}} - \sum_{k=1}^{r_\infty} A_{\infty,-k} x^{k-1}, \quad (1.1)$$

having regular or irregular singularities of arbitrary rank.

The monodromy data to be preserved is

- (i) Stokes multipliers $S_j^{(\nu)}$ ($j = 1, \dots, 2r_\nu$),
- (ii) connection matrices $C^{(\nu)}$,
- (iii) “exponents of formal monodromy” $T_0^{(\nu)}$.

Miwa-Jimbo found a deformation equation as a necessary and sufficient condition for the monodromy data to be invariant for deformation parameters and defined the τ function for the deformation equation.

We will explain the relationship between the τ function and the θ function. In [4], Miwa-Jimbo expressed the τ function with the θ function by moving the argument z of the θ function. The monodromy preserving deformation has irregular singularities. The Riemann surface is a ramified covering of $\mathbb{CP}^1$ with the covering degree m and its genus is g . Especially the case of genus one is written in [2], which we will describe in Appendix.

In [6], Kitaev-Korotkin calculated the τ function with the θ function by moving the period Ω of the θ function. They consider monodromy preserving deformation of a second order Fuchsian equation with $2g + 2$ regular singular points. In this case the τ function is represented by the theta null value of a hyperelliptic curve with genus g . In case $g = 1$, Kitaev-Korotkin's monodromy preserving deformation is equivalent to Picard's solution of the sixth Painlevé equation by the Bäcklund transformation. The aim of this paper is to unify Miwa-Jimbo's result and Kitaev-Korotkin's result in the elliptic case.

It is well known that the elliptic θ function satisfies the heat equation, whose solution space is infinite dimensional. The equation (5.4) of isomonodromic deformation is solved by the τ function. The elliptic θ function also satisfies the nonlinear equation (5.4), whose solution space is finite dimensional.

We will explain this paper in detail. In section 2, we show some definitions and facts of elliptic functions, which is necessary to construct the τ -function.

In section 3, we will study the following ordinary differential equation:

$$\frac{dY}{dx} = \left(\frac{B_{-1}}{(x-a)^2} + \frac{B_0}{x-a} + \sum_{\nu=1}^3 \frac{A_\nu}{x-e_\nu} \right) Y(x), \quad (1.2)$$

and define the τ -function.

In section 4, we construct a fundamental solution of (1.2) and calculate the following monodromy data:

- (i) Stokes multipliers around $x = a$ $S_1^a = S_2^a = 1$,
- (ii) connection matrices around $x = e_\nu$ $C^{(\nu)} (\nu = 1, 2, 3, \infty)$,
- (iii) the exponents of formal monodromy $T_0^a = 0, T_0^{(\nu)} = \text{diag}(-\frac{1}{4}, \frac{1}{4}) (\nu = 1, 2, 3, \infty)$,

where e_∞ is ∞ .

The behavior of the fundamental solution near the singular points is as follows

$$Y(x) = (1 + O(x-a)) \exp T^{(a)}(x), \quad (1.3)$$

$$Y(x) = G^{(\nu)}(1 + O(x-e_\nu)) \exp T^{(\nu)}(x) \quad (\nu = 1, 2, 3), \quad (1.4)$$

$$Y(x) = G^{(\infty)}(1 + O(\frac{1}{x})) \exp T^{(\infty)}(x) \quad (1.5)$$

where

$$T^{(a)}(x) = \begin{pmatrix} \frac{\wp'(\alpha)t}{2} & \\ & -\frac{\wp'(\alpha)t}{2} \end{pmatrix} \frac{(x-a)^{-1}}{-1} + T_0^{(a)} \log(x-a), \quad (1.6)$$

$$T^{(\nu)}(x) = T_0^{(\nu)} \log(x - e_\nu) \quad (\nu = 1, 2, 3), \quad (1.7)$$

$$T^{(\infty)}(x) = T_0^{(\infty)} \log\left(\frac{1}{x}\right). \quad (1.8)$$

In section 5, we concretely calculate the monodromy preserving deformation from the fundamental solution and compute the coefficients, B_{-1}, B_0, A_ν ($\nu = 1, 2, 3$).

In section 6, we calculate the τ -function. Section 6 consists of three subsections. Subsection 6.1 is devoted to H_t , the Hamiltonian on the deformation parameter t . Subsection 6.2, 6.3 is about H_ν ($\nu = 1, 2, 3$), the Hamiltonians on the deformation parameters e_ν ($\nu = 1, 2, 3$). In subsection 6.2, we show some facts about elliptic functions in order to calculate the Hamiltonians H_ν ($\nu = 1, 2, 3$). In subsection 6.3, we calculate H_ν ($\nu = 1, 2, 3$) and the τ -function. Our main theorem is as follows:

Theorem 1.1. *For the monodromy preserving deformation (1.2), the τ function is*

$$\tau(e_1, e_2, e_3, t) = \theta[p, q] \left(\frac{t}{\omega_1}; \Omega \right) \omega_1^{-\frac{1}{2}} \prod_{1 \leq \nu < \mu \leq 3} (e_\nu - e_\mu)^{-\frac{1}{8}} \exp\left(\frac{\eta_1 t^2}{2\omega_1}\right) \exp\left(\frac{t^2}{4} f(e_1, e_2, e_3)\right),$$

where

$$f(e_1, e_2, e_3) = -a + \frac{1}{3} \sum_{\nu=1}^3 e_\nu + \frac{1}{2} \prod_{\nu=1}^3 (a - e_\nu) \sum_{\mu=1}^3 \frac{1}{(a - e_\mu)^2}.$$

Acknowledgments. The author thanks Professor Yousuke Ohyaama for the careful guidance. This work is partially done by the author in the Isaac Newton Institute for Mathematical sciences, Cambridge University. The author is grateful to Professor D. Korotkin for a fruitful discussion.

2 Elliptic Function Theory

We will review the elliptic function theory to fix our notation. See [5]. We will consider the elliptic curve E defined by the equation

$$y^2 = 4(x - e_1)(x - e_2)(x - e_3)$$




Figure 1: Branch cuts and canonical basis of cycles on the elliptic curve, E . Continuous and dashed paths lie on the first and second sheet of E , respectively.

with arbitrary constants $e_i \in \mathbb{C}$ ($e_i \neq e_j$), where we do not assume that $e_1 + e_2 + e_3 = 0$. We set the new coordinate:

$$\tilde{x} = x - \frac{1}{3} \sum_{i=1}^3 e_i, \quad \tilde{e}_j = e_j - \frac{1}{3} \sum_{i=1}^3 e_i \quad \text{for } j = 1, 2, 3$$

and define the Abel map in the following way:

$$u := \int_{\infty}^x \frac{dx}{\sqrt{4(x - e_1)(x - e_2)(x - e_3)}} = \int_{\infty}^{\tilde{x}} \frac{d\tilde{x}}{\sqrt{4(\tilde{x} - \tilde{e}_1)(\tilde{x} - \tilde{e}_2)(\tilde{x} - \tilde{e}_3)}}. \quad (2.1)$$

Let us define periods ω_1, ω_2 by

$$\omega_1 = \int_{\gamma} \frac{dx}{y}, \quad \omega_2 = \int_{\delta} \frac{dx}{y}. \quad (2.2)$$

From the bilinear relation of Riemann, we get

$$\Omega := \frac{\omega_2}{\omega_1}, \quad \Im(\Omega) > 0. \quad (2.3)$$

$\tilde{x} = \tilde{x}(u)$, the inverse function of the Abel map, can be expressed with the Weierstrass $\wp$ -function:

$$x - \frac{1}{3} \sum_{i=1}^3 e_i = \tilde{x}(u) = \wp(u; \omega_1, \omega_2). \quad (2.4)$$

$\wp(u)$ satisfies the following differential equations:

$$\wp'(u)^2 = 4(\wp(u) - \tilde{e}_1)(\wp(u) - \tilde{e}_2)(\wp(u) - \tilde{e}_3) \quad (2.5)$$

$$= 4\wp(u)^3 - g_2\wp(u) - g_3 \quad (2.6)$$

We have

$$\sigma(u; \omega_1, \omega_2) = \exp\left(\frac{\eta_1}{2\omega_1}u^2\right) \frac{\omega_1}{\theta'_{11}} \theta_{11}\left(\frac{u}{2\omega_1}; \Omega\right). \quad (2.7)$$

We set the σ function with characteristic $p, q \in \mathbb{C}$ in the following way:

$$\sigma[p, q](u) := \exp\left(\frac{\eta_1}{2\omega_1}u^2\right) \frac{\omega_1}{\theta'_{11}} \theta[p, q]\left(\frac{u}{\omega_1}; \Omega\right). \quad (2.8)$$

$\sigma[p, q](u)$ has the following periodicity properties:

$$\sigma[p, q](u + \omega_1) = \exp(2\pi i p) \exp \eta_1 \left(u + \frac{\omega_1}{2}\right) \sigma[p, q](u), \quad (2.9)$$

$$\sigma[p, q](u + \omega_2) = \exp(-2\pi i q) \exp \eta_2 \left(u + \frac{\omega_2}{2}\right) \sigma[p, q](u). \quad (2.10)$$

3 The Schlesinger System

We will study the following ordinary differential equation:

$$\frac{dY}{dx} = \left(\frac{B_{-1}}{(x-a)^2} + \frac{B_0}{x-a} + \sum_{\nu=1}^3 \frac{A_\nu}{x-e_\nu} \right) Y(x), \quad (3.1)$$

where $A_1, A_2, A_3, B_{-1}, B_0 \in sl(2, \mathbb{C})$ are independent of x .

The monodromy data of (3.1) is as follows:

- (i) Stokes multipliers around $x = a$ $S_1^a = S_2^a$;
- (ii) connection matrices around ∞, e_ν $C^\infty, C^{(\nu)}$ ($\nu = 1, 2, 3$);
- (iii) the exponents of formal monodromy $T_0^a, T_0^{(\infty)}, T_0^{(\nu)}$ ($\nu = 1, 2, 3$).

In the next section, we will get convergent series around $x = a, \infty, e_\nu$ ($\nu = 1, 2, 3$) in the following way:

$$Y(x) = (1 + O(x-a)) \exp T^{(a)}(x) = \hat{Y}^{(a)}(x) \exp T^{(a)}(x), \quad (3.2)$$

$$Y(x) = G^{(\infty)}(1 + O(\frac{1}{x})) \exp T^{(\infty)}(x) = G^{(\infty)} \hat{Y}^{(\infty)}(x) \exp T^{(\infty)}(x), \quad (3.3)$$

$$Y(x) = G^{(\nu)}(1 + O(x-e_\nu)) \exp T^{(\nu)}(x) = G^{(\nu)} \hat{Y}^{(\nu)}(x) \exp T_0^{(\nu)}(x), \quad (\nu = 1, 2, 3). \quad (3.4)$$

where

$$T^{(a)}(x) = \begin{pmatrix} \frac{\wp'(\alpha)t}{2} & \\ & -\frac{\wp'(\alpha)t}{2} \end{pmatrix} \frac{(x-a)^{-1}}{-1} + T_0^{(a)} \log(x-a), \quad (3.5)$$

$$T^{(\infty)}(x) = T_0^\infty \log\left(\frac{1}{x}\right) \quad (3.6)$$

$$T^{(\nu)}(x) = T_0^{(\nu)} \log(x - e_\nu) \quad (\nu = 1, 2, 3). \quad (3.7)$$

For the deformation parameters t, e_1, e_2, e_3 , we set the following closed 1-form

$$\omega = \omega_a + \omega_{e_1} + \omega_{e_2} + \omega_{e_3} \quad (3.8)$$

$$= H_t dt + H_1 de_1 + H_2 de_2 + H_3 de_3, \quad (3.9)$$

where

$$\omega_a = -\text{Res}_{x=a} \text{tr} \hat{Y}^{(a)}(x)^{-1} \frac{\partial \hat{Y}^{(a)}}{\partial x}(x) dT^{(a)}(x), \quad (3.10)$$

$$\omega_{e_\nu} = -\text{Res}_{x=e_\nu} \text{tr} \hat{Y}^{(\nu)}(x)^{-1} \frac{\partial \hat{Y}^{(\nu)}}{\partial x}(x) dT^{(\nu)}(x) \quad (\nu = 1, 2, 3), \quad (3.11)$$

and d is the exterior differentiation with respect to the deformation parameters t, e_1, e_2, e_3 . Especially, we can write

$$\omega_{e_\nu} = \left[\text{Res}_{x=e_\nu} \frac{1}{2} \text{tr} \left(\frac{dY}{dx} Y^{-1} \right)^2 \right] de_\nu. \quad (3.12)$$

From the closed 1-form ω , we define the τ -function in the following way:

$$\omega := d \log \tau(e_1, e_2, e_3, t). \quad (3.13)$$

4 The Riemann-Hilbert Problem for Special Parameters

In this section, we will concretely construct a 2×2 matrix valued function $Y(x)$, whose monodromy data, (i) Stokes multipliers, (ii) connection matrices, (iii) exponents of formal monodromies, is independent of the deformation parameters t, e_1, e_2, e_3 .

We denote the Abel map $u = u(x)$ by $u = u(P)$, because we regard it as a function of the Riemann surface E defined by

$$y^2 = 4(x - e_1)(x - e_2)(x - e_3). \quad (4.1)$$




Figure 2: Generators of $\pi_1(\mathbb{CP}^1 \setminus \{\infty, e_1, e_2, e_3\}, x_0)$

We set the involution of E in the following way:

$$* : (x, y) \longrightarrow (x, -y). \quad (4.2)$$

We define the 2×2 matrix valued function $\Phi(P)$ with the σ function and the ζ function in the following way:

$$\Phi(P) = \begin{pmatrix} \varphi(P) & \varphi(P^*) \\ \psi(P) & \psi(P^*) \end{pmatrix}, \quad (4.3)$$

where

$$\begin{aligned} \varphi(P) &= \sigma[p, q](u + u_\varphi + t)\sigma(u - u_\varphi) \exp -\frac{t}{2}\{\zeta(u - \alpha) + \zeta(u + \alpha)\} \\ \psi(P) &= \sigma[p, q](u + u_\psi + t)\sigma(u - u_\psi) \exp -\frac{t}{2}\{\zeta(u - \alpha) + \zeta(u + \alpha)\} \\ u_\varphi &= u(P_\varphi), u_\psi = u(P_\psi), \alpha = u(a), \end{aligned}$$

with arbitrary $P_\varphi, P_\psi \in E$.

Proposition 4.1. *The function $\Phi(P)$ is holomorphic and invertible outside of the branch points ∞, e_1, e_2, e_3 and transforms as follows with respect to the tracing along the basic cycles γ, δ :*

$$\begin{aligned} T_\gamma[\Phi(P)] &= \Phi(P) \begin{pmatrix} \exp\{\pi i(2p+1)\} & \\ & \exp\{-\pi i(2p+1)\} \end{pmatrix} \exp\{\eta_1(2u + \omega_1)\} \\ T_\delta[\Phi(P)] &= \Phi(P) \begin{pmatrix} \exp\{-\pi i(2q+1)\} & \\ & \exp\{\pi i(2q+1)\} \end{pmatrix} \exp\{\eta_2(2u + \omega_2)\} \end{aligned}$$

where by T_l we denote the operator of analytic continuation along the contour l .

Proof. By using the periodicity (2.9) and (2.10), we get

$$\begin{aligned} T_\gamma [\Phi(P)] &= \Phi(P) \begin{pmatrix} \exp\{\pi i(2p+1)\} & \\ & \exp\{-\pi i(2p+1)\} \end{pmatrix} \exp\{\eta_1(2u+\omega_1)\} \\ T_\delta [\Phi(P)] &= \Phi(P) \begin{pmatrix} \exp\{-\pi i(2q+1)\} & \\ & \exp\{\pi i(2q+1)\} \end{pmatrix} \exp\{\eta_2(2u+\omega_2)\}. \end{aligned}$$

Then we obtain

$$\begin{aligned} T_\gamma [\det \Phi(P)] &= \exp 2\eta_1(2u+\omega_1) \det \Phi(P) \\ T_\delta [\det \Phi(P)] &= \exp 2\eta_2(2u+\omega_2) \det \Phi(P). \end{aligned}$$

By integrating $\frac{d(\det \Phi(P))}{\det \Phi(P)}$ along the fundamental polygon of E and using the Legendre relation, we get

$$\frac{1}{2\pi i} \oint_{\partial \hat{E}} \frac{d(\det \Phi(P))}{\det \Phi(P)} = \frac{1}{2\pi i} 4(\eta_1\omega_2 - \eta_2\omega_1) = 4. \quad (4.4)$$

Therefore, four zeros of $\det \Phi(P)$ on E are branch points, ∞, e_1, e_2, e_3 . We proved the proposition. $\square$

We will normalize $\Phi(P)$ at $x = a$. We can easily show the local behavior of the Abel map.

Lemma 4.2.

$$u - \alpha = \frac{1}{\wp'(\alpha)}(x - a) - \frac{\wp''(\alpha)}{2\wp'''(\alpha)}(x - a)^2 + \left(\frac{\wp''(\alpha)^2}{2\wp'(\alpha)^5} - \frac{\wp'''(\alpha)}{6\wp'(\alpha)^4} \right) (x - a)^3 + \dots \quad (4.5)$$

From Lemma 4.2, $\Phi(x)$ can be developed near $x = a$ in the following way:

$$\Phi(x) = (G^{(a)} + O(x - a)) \exp \begin{pmatrix} -\frac{\wp'(\alpha)t}{2(x-a)} & \\ & \frac{\wp'(\alpha)t}{2(x-a)} \end{pmatrix},$$

where $G^{(a)}$ is a 2×2 matrix with the matrix elements:

$$\begin{aligned} (G^{(a)})_{11} &= \sigma[p, q](\alpha + u_\varphi + t)\sigma(\alpha - u_\varphi) \exp \left\{ -\frac{t}{2} \left(\frac{\wp''(\alpha)}{2\wp'(\alpha)} + \zeta(2\alpha) \right) \right\}, \\ (G^{(a)})_{21} &= \sigma[p, q](\alpha + u_\psi + t)\sigma(\alpha - u_\psi) \exp \left\{ -\frac{t}{2} \left(\frac{\wp''(\alpha)}{2\wp'(\alpha)} + \zeta(2\alpha) \right) \right\}, \\ (G^{(a)})_{12} &= \sigma[p, q](-\alpha + u_\varphi + t)\sigma(-\alpha - u_\varphi) \exp \left\{ \frac{t}{2} \left(\frac{\wp''(\alpha)}{2\wp'(\alpha)} + \zeta(2\alpha) \right) \right\}, \\ (G^{(a)})_{22} &= \sigma[p, q](-\alpha - u_\psi + t)\sigma(-\alpha - u_\psi) \exp \left\{ \frac{t}{2} \left(\frac{\wp''(\alpha)}{2\wp'(\alpha)} + \zeta(2\alpha) \right) \right\}. \end{aligned}$$

From Proposition 4.1,

$$\det G^{(a)} = \det \Phi(a) \neq 0.$$

We define a matrix valued function $Y(x)$ in the following way:

$$Y(P) = \frac{\sqrt{\det \Phi(a)}}{\sqrt{\det \Phi(P)}} (G^{(a)})^{-1} \Phi(P), \quad (4.6)$$

By the normalization near $x = a$, we obtain the following lemma:

Lemma 4.3.

$$Y(x) = \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + O(x - a) \right) \exp T^{(a)}(x)$$

$$T^{(a)}(x) = T_{-1}^{(a)} \frac{(x - a)^{-1}}{-1}, \quad T_{-1}^{(a)} = \begin{pmatrix} \frac{\wp'(\alpha)t}{2} & \\ & \frac{-\wp'(\alpha)t}{2} \end{pmatrix}.$$

Especially, if $u_\varphi = \alpha, u_\psi = -\alpha$,

$$Y(x) = \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + Y_1^{(a)}(x - a) + \cdots \right) \exp T^{(a)}(x)$$

$$\begin{aligned} \left(Y_1^{(a)} \right)_{11} &= \frac{1}{\wp'(\alpha)} \left\{ \frac{\sigma[p, q]'(t)}{\sigma[p, q](t)} - \frac{t}{2} \left(4\wp(\alpha) - \frac{1}{2} \left(\frac{\wp''(\alpha)}{\wp'(\alpha)} \right)^2 \right) \right\} \\ \left(Y_1^{(a)} \right)_{21} &= \frac{\sigma[p, q](2\alpha + t)}{\sigma[p, q](t)\sigma[p, q](2\alpha)\wp'(\alpha)} \exp \left\{ -t \left(\frac{\wp''(\alpha)}{2\wp'(\alpha)} + \zeta(2\alpha) \right) \right\} \\ \left(Y_1^{(a)} \right)_{12} &= \frac{-\sigma[p, q](-2\alpha + t)}{\sigma[p, q](t)\sigma[p, q](2\alpha)\wp'(\alpha)} \exp \left\{ t \left(\frac{\wp''(\alpha)}{2\wp'(\alpha)} + \zeta(2\alpha) \right) \right\} \\ \left(Y_1^{(a)} \right)_{22} &= \frac{-1}{\wp'(\alpha)} \left\{ \frac{\sigma[p, q]'(t)}{\sigma[p, q](t)} - \frac{t}{2} \left(4\wp(\alpha) - \frac{1}{2} \left(\frac{\wp''(\alpha)}{\wp'(\alpha)} \right)^2 \right) \right\}. \end{aligned}$$

$Y(x)$ has the following monodromy and Stokes matrices.

Theorem 4.4. *For $\nu = \infty, 1, 2, 3$, we set*

$$M_\nu = \begin{pmatrix} 0 & m_\nu \\ -m_\nu^{-1} & 0 \end{pmatrix}. \quad (4.7)$$

The matrix elements corresponding to the contour l_ν ($\nu = \infty, 1, 2, 3$) are given by the expressions

$$\begin{aligned} m_\infty &= -i, \quad m_1 = i \exp(-2\pi i p) \\ m_2 &= -i \exp 2\pi i(-p + q) \\ m_3 &= i \exp(2\pi i q). \end{aligned}$$

Stokes matrices are as follows

$$S_1 = S_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Proof. From Proposition 4.1, we note that the branch points ∞, e_1, e_2, e_3 on E are zero points of $\det \Phi(P)$ of the first order. According to Proposition 4.1, we obtain

$$\begin{aligned} T_\gamma [Y(x)] &= Y(x) M_3 M_2 \\ &= Y(x) \begin{pmatrix} \exp\{2\pi ip\} & \\ & \exp\{-2\pi ip\} \end{pmatrix}, \end{aligned}$$

$$\begin{aligned} T_{\delta^{-1}} [Y(x)] &= Y(x) M_2 M_1 \\ &= Y(x) \begin{pmatrix} \exp\{2\pi iq\} & \\ & \exp\{-2\pi iq\} \end{pmatrix}. \end{aligned}$$

By the involution $*$, we get

$$Y(x) M_\infty = Y(x) \begin{pmatrix} & i \\ i & \end{pmatrix}, \text{ or } = Y(x) \begin{pmatrix} & -i \\ -i & \end{pmatrix}.$$

We set

$$M_\infty = \begin{pmatrix} & m_\infty \\ -m_\infty^{-1} & \end{pmatrix} = \begin{pmatrix} & -i \\ -i & \end{pmatrix}.$$

From the property of the monodromy group, we obtain

$$M_3 M_2 M_1 = M_\infty^{-1}. \tag{4.8}$$

From

$$\begin{aligned} M_3 M_2 &= \text{diag}(\exp\{2\pi ip\}, \exp\{-2\pi ip\}), \\ M_2 M_1 &= \text{diag}(\exp\{2\pi iq\}, \exp\{-2\pi iq\}), \end{aligned}$$

we get

$$\begin{aligned} M_1 &= \begin{pmatrix} & -m_\infty \exp\{-2\pi ip\} \\ m_\infty^{-1} \exp\{2\pi ip\} & \end{pmatrix}, \\ M_3 &= \begin{pmatrix} & -m_\infty \exp\{2\pi iq\} \\ m_\infty^{-1} \exp\{-2\pi iq\} & \end{pmatrix}. \end{aligned}$$

Therefore, we get

$$M_2 = \begin{pmatrix} & m_\infty \exp\{2\pi i(q-p)\} \\ -m_\infty^{-1} \exp\{2\pi i(p-q)\} & \end{pmatrix},$$

from (4.8).

From Lemma 4.3, the Stokes matrices are

$$S_1 = S_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

□

We can describe the monodromy data of $Y(x)$ in the following way.

Corollary 4.5. *$Y(x)$ has the following monodromy data:*

(i) *Stokes multipliers $S_1^a = S_2^a = 1$,*

(ii) *connection matrices $C^{(\nu)} = \frac{1}{\sqrt{2im_\nu}} \begin{pmatrix} i & -m_\nu \\ i & m_\nu \end{pmatrix}$ ($\nu = \infty, 1, 2, 3$),*

(iii) *formal monodromies $T_0^{(\nu)} = \text{diag}(-\frac{1}{4}, \frac{1}{4})$ ($\nu = \infty, 1, 2, 3$).*

Especially, the behaviors of $Y(x)$ near e_ν ($\nu = \infty, 1, 2, 3$) are

$$\begin{aligned} Y(x) &= G^{(\nu)}(1 + O(x_\nu)) \exp T^{(\nu)}(x) C^{(\nu)} \\ T^{(\nu)}(x) &= T_0^{(\nu)}(x) \log x_\nu \\ x_\infty &= x^{-1}, \quad x_\nu = x - e_\nu \end{aligned}$$

Proof. (i) is clear. (ii) and (iii) can be obtained by diagonalizing the monodromy matrices M_ν ($\nu = \infty, 1, 2, 3$). □

This corollary means that the monodromy data of $Y(x)$ is independent of the deformation parameters t, e_1, e_2, e_3 .

5 Monodromy Preserving Deformation

In this section, we will prove that $Y(x)$ satisfies an ordinary differential equation of the following type:

$$\frac{dY}{dx} = \left(\frac{B_{-1}}{(x-a)^2} + \frac{B_0}{x-a} + \sum_{\nu=1}^3 \frac{A_\nu}{x-e_\nu} \right) Y(x),$$

and concretely determine the coefficients B_{-1}, B_0, A_ν ($\nu = 1, 2, 3$).

Firstly, we define symbols in order to describe the monodromy preserving deformation. For $\nu = 1, 2, 3$, we set

$$\begin{aligned} D^{(\nu)}(u) &= \frac{2m_\nu}{m_\infty} \varphi(u) \psi(u) \left(\frac{d}{du} \log \varphi(u) - \frac{d}{du} \log \psi(u) \right), \\ \tilde{\omega}_1 &= \frac{\omega_1}{2}, \tilde{\omega}_2 = \frac{\omega_1 + \omega_2}{2}, \tilde{\omega}_3 = \frac{\omega_2}{2}, \\ \tilde{\eta}_1 &= \eta_1, \tilde{\eta}_2 = \eta_1 + \eta_2, \tilde{\eta}_3 = \eta_2. \end{aligned}$$

We note that the monodromy matrices and the Stoke matrices do not have the parameters P_φ, P_ϕ in Theorem 4.4. Then in this section, we set $u_\varphi = \alpha$, $u_\phi = -\alpha$ from the uniqueness of the Riemann-Hilbert problem.

Theorem 5.1. *$Y(x)$ satisfies the following ordinary differential equation:*

$$\frac{dY}{dx} = \left(\frac{B_{-1}}{(x-a)^2} + \frac{B_0}{x-a} + \sum_{\nu=1}^3 \frac{A_\nu}{x-e_\nu} \right) Y(x), \quad (5.1)$$

where

$$\begin{aligned} B_{-1} &= \text{diag} \left(\frac{\wp'(\alpha)t}{2}, -\frac{\wp'(\alpha)t}{2} \right) \\ B_0 &= \text{diag}(0, 0) \\ A_\nu &= G^{(\nu)} \text{diag} \left(-\frac{1}{4}, \frac{1}{4} \right) (G^{(\nu)})^{-1}, \quad (\nu = 1, 2, 3), \\ G^{(\nu)} &= \begin{pmatrix} 0 & \exp \left\{ \frac{t}{2} \left(\frac{\wp''(\alpha)}{2\wp'(\alpha)} + \zeta(2\alpha) \right) \right\} \\ -\exp \left\{ -\frac{t}{2} \left(\frac{\wp''(\alpha)}{2\wp'(\alpha)} + \zeta(2\alpha) \right) \right\} & 0 \end{pmatrix} \\ &\quad \times \frac{\sqrt{2m_\nu}}{\sqrt{D^{(\nu)}(\tilde{\omega}_\nu)}i} \begin{pmatrix} \varphi(\tilde{\omega}_\nu) & \varphi(\tilde{\omega}_\nu) \left(\frac{d}{du} \log \varphi(\tilde{\omega}_\nu) - \tilde{\eta}_\nu \right) \\ \psi(\tilde{\omega}_\nu) & \psi(\tilde{\omega}_\nu) \left(\frac{d}{du} \log \psi(\tilde{\omega}_\nu) - \tilde{\eta}_\nu \right) \end{pmatrix} \begin{pmatrix} \left(\frac{\wp''(\tilde{\omega}_\nu)}{2} \right)^{\frac{1}{4}} & \\ & \left(\frac{\wp''(\tilde{\omega}_\nu)}{2} \right)^{-\frac{1}{4}} \end{pmatrix}. \end{aligned}$$

The behavior near ∞ is

$$Y(x) = G^{(\infty)} \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + O(x^{-1}) \right) \exp T^{(\infty)}(x),$$

where

$$G^{(\infty)} = \begin{pmatrix} 0 & \exp \left\{ \frac{t}{2} \left(\frac{\wp''(\alpha)}{2\wp'(\alpha)} + \zeta(2\alpha) \right) \right\} \\ \exp \left\{ -\frac{t}{2} \left(\frac{\wp''(\alpha)}{2\wp'(\alpha)} + \zeta(2\alpha) \right) \right\} & 0 \end{pmatrix} \\ \times \frac{1}{\left(\frac{d}{du} \log \varphi(0) - \frac{d}{du} \log \psi(0) \right)^{\frac{1}{2}}} \begin{pmatrix} -i\varphi(0) & i\varphi'(0) \\ -i\psi(0) & i\psi'(0) \end{pmatrix}.$$

Proof. From Lemma 4.3, we get

$$Y'(x)Y^{-1}(x) = \frac{1}{(x-a)^2} \begin{pmatrix} \frac{\wp'(\alpha)t}{2} & \\ & -\frac{\wp'(\alpha)t}{2} \end{pmatrix} + \text{regular part} \\ := \frac{B_{-1}}{(x-a)^2} + \frac{B_0}{x-a} + \text{regular part}.$$

Therefore B_{-1}, B_0 are determined as in Theorem 5.1

We represent the monodromy matrices M_ν ($\nu = 1, 2, 3$) in the following way:

$$M_\nu = (C^{(\nu)})^{-1} \begin{pmatrix} \exp\{-\frac{2\pi i}{4}\} & \\ & \exp\{\frac{2\pi i}{4}\} \end{pmatrix} C^{(\nu)},$$

where $C^{(\nu)}$ is the connection matrix. We set $G^{(\nu)}$ ($\nu = 1, 2, 3$) in the following way;

$$Y(x) = G^{(\nu)} \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + O(x - e_\nu) \right) (x - e_\nu) \begin{pmatrix} -\frac{1}{4} & \\ & \frac{1}{4} \end{pmatrix} C^{(\nu)}. \quad (5.2)$$

In order to calculate $G^{(\nu)}$, we compute the constant term of

$$Y(x)(C^{(\nu)})^{-1}(x - e_\nu) \begin{pmatrix} \frac{1}{4} & \\ & -\frac{1}{4} \end{pmatrix} = Y(x) \frac{1}{\sqrt{2im_\nu}} \begin{pmatrix} m_\nu & m_\nu \\ -i & i \end{pmatrix} (x - e_\nu) \begin{pmatrix} \frac{1}{4} & \\ & -\frac{1}{4} \end{pmatrix} \\ = \frac{\sqrt{\det \Phi(a)}}{\sqrt{\det \Phi(u)}} (G^{(a)})^{-1} \frac{1}{\sqrt{2im_\nu}} \begin{pmatrix} m_\nu \varphi(u) - i\varphi(-u) & m_\nu \varphi(u) + i\varphi(-u) \\ m_\nu \psi(u) - i\psi(-u) & m_\nu \psi(u) + i\psi(-u) \end{pmatrix} (x - e_\nu) \begin{pmatrix} \frac{1}{4} & \\ & -\frac{1}{4} \end{pmatrix}.$$

By setting $u_\varphi = \alpha, u_\psi = -\alpha$ in $G^{(a)}$,

$$\sqrt{\det \Phi(a)} (G^{(a)})^{-1} = \begin{pmatrix} 0 & \exp \left\{ \frac{t}{2} \left(\frac{\wp''(\alpha)}{2\wp'(\alpha)} + \zeta(2\alpha) \right) \right\} \\ -\exp \left\{ -\frac{t}{2} \left(\frac{\wp''(\alpha)}{2\wp'(\alpha)} + \zeta(2\alpha) \right) \right\} & 0 \end{pmatrix}. \quad (5.3)$$

By direct computation, we get

$$\begin{aligned}
\det \Phi(u) &= (u - \tilde{\omega}_\nu) (\varphi'(\tilde{\omega}_\nu)\psi(-\tilde{\omega}_\nu) - \varphi(\tilde{\omega}_\nu)\psi'(-\tilde{\omega}_\nu) + \varphi'(-\tilde{\omega}_\nu)\psi(\tilde{\omega}_\nu) - \varphi(-\tilde{\omega}_\nu)\psi'(\tilde{\omega}_\nu)) + \dots \\
&= (u - \tilde{\omega}_\nu) \left\{ \varphi(\tilde{\omega}_\nu)\psi(-\tilde{\omega}_\nu) \left(\frac{\varphi'(\tilde{\omega}_\nu)}{\varphi(\tilde{\omega}_\nu)} - \frac{\psi'(-\tilde{\omega}_\nu)}{\psi(-\tilde{\omega}_\nu)} \right) \right. \\
&\quad \left. + \varphi(-\tilde{\omega}_\nu)\psi(\tilde{\omega}_\nu) \left(\frac{\varphi'(-\tilde{\omega}_\nu)}{\varphi(-\tilde{\omega}_\nu)} - \frac{\psi'(\tilde{\omega}_\nu)}{\psi(\tilde{\omega}_\nu)} \right) \right\} + \dots.
\end{aligned}$$

From (2.9), (2.10),

$$\begin{aligned}
\det \Phi(u) &= (u - \tilde{\omega}_\nu) \frac{m_\nu}{m_\infty} \varphi(\tilde{\omega}_\nu)\psi(\tilde{\omega}_\nu) \times 2 \left(\frac{d}{du} \log \varphi(\tilde{\omega}_\nu) - \frac{d}{du} \log \psi(\tilde{\omega}_\nu) \right) + O((u - \tilde{\omega}_\nu)^2) \\
&= (u - \tilde{\omega}_\nu) D^{(\nu)}(\tilde{\omega}_\nu) + O((u - \tilde{\omega}_\nu)^2) \\
&= (x - e_\nu)^{\frac{1}{2}} \left(\frac{\wp''(\tilde{\omega}_\nu)}{2} \right)^{-\frac{1}{2}} D^{(\nu)}(\tilde{\omega}_\nu) + O((\sqrt{x - e_\nu})^2)
\end{aligned}$$

From (2.9), (2.10), we note that

$$\begin{aligned}
m_\nu \varphi(\tilde{\omega}_\nu) + i \varphi(-\tilde{\omega}_\nu) &= 0, \\
m_\nu \psi(\tilde{\omega}_\nu) + i \psi(-\tilde{\omega}_\nu) &= 0.
\end{aligned}$$

Therefore,

$$\begin{aligned}
G^{(\nu)} &= Y(x) (C^{(\nu)})^{-1} (x - e_\nu) \begin{pmatrix} \frac{1}{4} & \\ & -\frac{1}{4} \end{pmatrix} \Bigg|_{x=e_\nu} \\
&= (\det \Phi(a))^{\frac{1}{2}} (G^{(a)})^{-1} (D^{(\nu)}(\tilde{\omega}_\nu))^{-\frac{1}{2}} \left(\frac{\wp''(\tilde{\omega}_\nu)}{2} \right)^{\frac{1}{4}} \\
&\quad \times \frac{1}{\sqrt{2im_\nu}} \begin{pmatrix} m_\nu \varphi(u) - i \varphi(-u) & \frac{m_\nu \varphi(u) + i \varphi(-u)}{\frac{u - \tilde{\omega}_\nu}{u - \tilde{\omega}_\nu}} \\ m_\nu \psi(u) - i \psi(-u) & \frac{m_\nu \psi(u) + i \psi(-u)}{\frac{u - \tilde{\omega}_\nu}{u - \tilde{\omega}_\nu}} \end{pmatrix} \begin{pmatrix} 1 & \\ & \frac{u - \tilde{\omega}_\nu}{(x - e_\nu)^{\frac{1}{2}}} \end{pmatrix} \Bigg|_{x=e_\nu} \\
&= (\det(a))^{\frac{1}{2}} (G^{(a)})^{-1} \frac{\sqrt{2m_\nu}}{\sqrt{D^{(\nu)}(\tilde{\omega}_\nu)i}} \begin{pmatrix} \varphi(\tilde{\omega}_\nu) & \varphi(\tilde{\omega}_\nu) \left(\frac{d}{du} \log \varphi(\tilde{\omega}_\nu) - \tilde{\eta}_\nu \right) \\ \psi(\tilde{\omega}_\nu) & \psi(\tilde{\omega}_\nu) \left(\frac{d}{du} \log \psi(\tilde{\omega}_\nu) - \tilde{\eta}_\nu \right) \end{pmatrix} \\
&\quad \times \begin{pmatrix} \left(\frac{\wp''(\tilde{\omega}_\nu)}{2} \right)^{\frac{1}{4}} & \\ & \left(\frac{\wp''(\tilde{\omega}_\nu)}{2} \right)^{-\frac{1}{4}} \end{pmatrix}.
\end{aligned}$$

From (5.2), we get

$$\begin{aligned} Y'(x)Y^{-1}(x) &= \frac{1}{(x - e_\nu)} G^{(\nu)} \begin{pmatrix} -\frac{1}{4} & \\ & \frac{1}{4} \end{pmatrix} (G^{(\nu)})^{-1} + \text{regular part} \\ &:= \frac{A_\nu}{(x - e_\nu)} + \text{regular part}. \end{aligned}$$

We set $G^{(\infty)}$ in the following way:

$$Y(x) = G^{(\infty)} (1 + O(x^{-1})) \exp T^{(\infty)}(x).$$

In the same way, we can compute $G^{(\infty)}$. □

We can get the following deformation equation from Theorem 5.1.

Corollary 5.2. *The deformation equation of the monodromy preserving deformation (5.1) is as follows. For $\nu = 1, 2, 3$,*

$$\begin{aligned} dA_\nu &= \sum_{\mu \neq \nu} \frac{[A_\mu, A_\nu]}{e_\nu - e_\mu} de_\nu + \sum_{\mu} \frac{[A_\mu, A_\nu]}{a - e_\mu} de_\mu \\ &\quad - \frac{[A_\nu, B_{-1}]}{(a - e_\nu)^2} de_\nu + \frac{[dT_{-1}^{(a)}, A_\nu]}{a - e_\nu} + [[dT_{-1}^{(a)}, Y_1^{(a)}], A_\nu], \end{aligned} \quad (5.4)$$

where d is the exterior differentiation with respect to the deformation parameters, t, e_1, e_2, e_3 and $T_{-1}^{(a)}, Y_1^{(a)}$ is defined in Lemma 4.3.

6 The tau-function for the Schlesinger System

In this section, we will calculate the τ -function for (5.1). This section consists of three subsections. Subsection 6.1 is devoted to the Hamiltonian H_t . Subsection 6.2, 6.3 is devoted to the Hamiltonian H_ν ($\nu = 1, 2, 3$). In subsection 6.2, we show some facts about elliptic functions. In subsection 6.3, we calculate H_ν and the τ -function.

6.1 The Hamiltonian at the Irregular Singular Point

In this subsection, we compute ω_a, H_t in the following way:

Proposition 6.1.

$$\begin{aligned} \omega_a &= \frac{\sigma[p, q]'(t)}{\sigma[p, q](t)} dt + \frac{\sigma[p, q]'(t)}{\sigma[p, q](t)} t \left(-\frac{de_1}{2(a - e_1)} - \frac{de_2}{2(a - e_2)} - \frac{de_3}{2(a - e_3)} \right) \\ &\quad + f(e_1, e_2, e_3) \left(\frac{tdt}{2} - \frac{t^2 de_1}{4(a - e_1)} - \frac{t^2 de_2}{4(a - e_2)} - \frac{t^2 de_3}{4(a - e_3)} \right), \end{aligned}$$

where

$$f(e_1, e_2, e_3) = -a + \frac{1}{3} \sum_{\nu=1}^3 e_\nu + \frac{1}{2} \prod_{\nu=1}^3 (a - e_\nu) \sum_{\mu=1}^3 \frac{1}{(a - e_\mu)^2}. \quad (6.1)$$

From ω_a , we get

$$\begin{aligned} H_t &= \frac{\sigma[p, q]'(t)}{\sigma[p, q](t)} + \frac{t}{2} f(e_1, e_2, e_3) \\ &= \frac{\partial}{\partial t} \left\{ \frac{\eta_1 t^2}{2\omega_1} \right\} + \frac{\partial}{\partial t} \left\{ \log \theta[p, q] \left(\frac{t}{\omega_1} \right) \right\} + \frac{\partial}{\partial t} \left\{ \frac{t^2}{4} f(e_1, e_2, e_3) \right\}. \end{aligned}$$

Proof. From the definition of ω_a ,

$$\omega_a = -\text{Res}_{x=a} \text{tr} \hat{Y}^{(a)}(x)^{-1} \frac{\partial \hat{Y}^{(a)}}{\partial x}(x) dT^{(a)}(x). \quad (6.2)$$

We set

$$\Pi(P) := -\frac{t}{2} \{ \zeta(u - \alpha) + \zeta(u + \alpha) \}. \quad (6.3)$$

From Lemma 4.2, we get

$$\begin{aligned} \Pi(P) &= -\frac{t}{2} \left\{ \frac{1}{u - \alpha} + \cdots + \zeta(u + \alpha) \right\} \\ &= -\frac{\wp'(\alpha)t}{2} \frac{1}{(x - a)} - \frac{t}{2} \left(\frac{\wp''(\alpha)}{2\wp'(\alpha)} + \zeta(2\alpha) \right) \\ &\quad - \frac{t}{2} \left(-\frac{\wp(2\alpha)}{\wp'(\alpha)} - \frac{\wp''(\alpha)^2}{4\wp'(\alpha)^3} + \frac{\wp'''(\alpha)}{6\wp'(\alpha)^2} \right) (x - a) + \cdots. \end{aligned}$$

We denote the regular part of $\Pi(P)$ around $x = a$ by $\hat{\Pi}(P)$:

$$\hat{\Pi}(P) := \Pi(P) - \left(-\frac{\wp'(\alpha)t}{2} \frac{1}{(x - a)} \right). \quad (6.4)$$

We set

$$\begin{aligned} \hat{\varphi}(P) &= \sigma[p, q](u + u_\varphi + t) \sigma(u - u_\varphi), \\ \hat{\psi}(P) &= \sigma[p, q](u + u_\psi + t) \sigma(u - u_\psi). \end{aligned}$$

Then we get

$$\begin{aligned}
Y(P) &= \frac{\sqrt{\det \Phi(a)}}{\sqrt{\det \Phi(P)}} (G^{(a)})^{-1} \Phi(P) \\
&= \frac{\sqrt{\det \Phi(a)}}{\sqrt{\det \Phi(P)}} (G^{(a)})^{-1} \begin{pmatrix} \hat{\varphi}(P) \exp(\hat{\Pi}(P)) & \hat{\varphi}(P^*) \exp(\hat{\Pi}(P^*)) \\ \hat{\psi}(P) \exp(\hat{\Pi}(P)) & \hat{\psi}(P^*) \exp(\hat{\Pi}(P^*)) \end{pmatrix} \\
&\quad \text{diag} \left(\exp \left\{ -\frac{\wp'(\alpha)t}{2} \frac{1}{x-a} \right\}, \exp \left\{ \frac{\wp'(\alpha)t}{2} \frac{1}{x-a} \right\} \right) \\
&:= \hat{Y}^{(a)}(x) \exp T^{(a)}(x).
\end{aligned}$$

According to the definition of ω_a ,

$$\omega_a = -\text{Res}_{x=a} \text{tr} \hat{Y}^{(a)}(x)^{-1} \frac{\partial}{\partial x} \hat{Y}^{(a)}(x) dT^{(a)}(x). \quad (6.5)$$

We set

$$A(x) = (G^{(a)})^{-1} \begin{pmatrix} \hat{\varphi}(u) \exp \hat{\Pi}(u) & \hat{\varphi}(-u) \exp \hat{\Pi}(-u) \\ \hat{\phi}(u) \exp \hat{\Pi}(u) & \hat{\phi}(-u) \exp \hat{\Pi}(-u) \end{pmatrix}. \quad (6.6)$$

By direct calculation, we get

$$-\text{tr} \hat{Y}^{(a)}(x)^{-1} \frac{\partial}{\partial x} \hat{Y}^{(a)}(x) dT^{(a)}(x) = \frac{d(\wp'(\alpha)t)}{2} \frac{1}{(x-a)} \text{tr} A^{-1}(x) A'(x) \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}. \quad (6.7)$$

We obtain

$$\begin{aligned}
&\text{tr} A^{-1}(x) A'(x) \begin{pmatrix} 1 & \\ & -1 \end{pmatrix} \\
&= \frac{1}{\det \Phi(u)} \left[\det \begin{pmatrix} \{\hat{\varphi}(u) \exp \hat{\Pi}(u)\}' & \hat{\varphi}(-u) \exp \hat{\Pi}(-u) \\ \{\hat{\psi}(u) \exp \hat{\Pi}(u)\}' & \hat{\psi}(-u) \exp \hat{\Pi}(-u) \end{pmatrix} \right. \\
&\quad \left. - \det \begin{pmatrix} \hat{\varphi}(u) \exp \hat{\Pi}(u) & \{\hat{\varphi}(-u) \exp \hat{\Pi}(-u)\}' \\ \hat{\psi}(u) \exp \hat{\Pi}(u) & \{\hat{\psi}(-u) \exp \hat{\Pi}(-u)\}' \end{pmatrix} \right],
\end{aligned}$$

where $'$ means the differentiation with respect to the variable x .

We normalized the matrix function $Y(x)$ around $x = a$ like Lemma 4.3 and proved that the monodromy group of $Y(x)$ is independent of P_φ, P_ψ in Theorem 4.4 and its corollary. Therefore, we can choose the parameters, P_φ, P_ψ at our disposal to simplify the calculation. Firstly, we multiply both the numerators and the denominators by $\frac{1}{x_\psi - x_\varphi}$. We take the limit $P_\psi \rightarrow P_\varphi$ and get

$$\hat{\psi}(P) = \frac{\partial \hat{\varphi}(P)}{\partial x_\varphi}. \quad (6.8)$$

Secondly, we multiply both the numerators and the denominators by $\frac{1}{x_\varphi - x}$. We take the limit $P_\varphi \rightarrow P$ and obtain

$$\operatorname{tr} A^{-1}(x)A'(x) \begin{pmatrix} 1 & \\ & -1 \end{pmatrix} = 2 \frac{1}{\sigma[p, q](t)} \lim_{P_\varphi \rightarrow P} \frac{\partial}{\partial x_\varphi} \sigma[p, q](-u + u_\varphi + t) + 2 \frac{\partial}{\partial x} \{\hat{\Pi}(P)\}. \quad (6.9)$$

According to the definition of ω_a ,

$$\begin{aligned} \omega_a &= \frac{d(\wp'(\alpha)t)}{2} \operatorname{tr} A^{-1}(x)A'(x) \begin{pmatrix} 1 & \\ & -1 \end{pmatrix} \Big|_{x=a} \\ &= \frac{\sigma[p, q]'(t)}{\sigma[p, q](t)} dt + \frac{\sigma[p, q]'(t)}{\sigma[p, q](t)} t \frac{d\wp'(\alpha)}{\wp'(\alpha)} \\ &\quad + \left(\frac{t}{2} dt + \frac{t^2}{2} \frac{d\wp'(\alpha)}{\wp'(\alpha)} \right) \left(\wp(2\alpha) + \frac{\wp''(\alpha)^2}{4\wp'(\alpha)^2} - \frac{\wp'''(\alpha)}{6\wp'(\alpha)} \right) \end{aligned}$$

where d is the exterior differentiation with respect to the deformation parameters, e_1, e_2, e_3, t .

In order to compute ω_a , we need some preparations about the $\wp$ -function. By direct calculation, we get

$$\frac{d\wp'(\alpha)}{\wp'(\alpha)} = -\frac{1}{2} \frac{1}{a - e_1} de_1 - \frac{1}{2} \frac{1}{a - e_2} de_2 - \frac{1}{2} \frac{1}{a - e_3} de_3. \quad (6.10)$$

We will use the addition theorem

$$\wp(2\alpha) = -2\wp(\alpha) + \frac{1}{4} \left(\frac{\wp''(\alpha)}{\wp'(\alpha)} \right)^2, \quad (6.11)$$

and the following equation:

$$\wp'''(\alpha) = 12\wp'(\alpha)\wp(\alpha). \quad (6.12)$$

From the relationship between the $\wp$ -function and the Abel-map, we obtain

$$\begin{aligned} \wp(\alpha) &= a - \frac{1}{3} \sum_{i=1}^3 e_i \\ \wp'(\alpha)^2 &= 4(a - e_1)(a - e_2)(a - e_3) \\ \wp''(\alpha) &= \sum_{i < j} 2(a - e_i)(a - e_j). \end{aligned}$$

By using the above equations, we get

$$\begin{aligned} \omega_a &= \frac{\sigma[p, q]'(t)}{\sigma[p, q](t)} dt + \frac{\sigma[p, q]'(t)}{\sigma[p, q](t)} t \left(-\frac{de_1}{2(a - e_1)} - \frac{de_2}{2(a - e_2)} - \frac{de_3}{2(a - e_3)} \right) \\ &\quad + f(e_1, e_2, e_3) \left(\frac{tdt}{2} - \frac{t^2 de_1}{4(a - e_1)} - \frac{t^2 de_2}{4(a - e_2)} - \frac{t^2 de_3}{4(a - e_3)} \right). \end{aligned}$$

□

6.2 Three Lemmas about Elliptic Functions

We show three lemmas about elliptic functions.

Lemma 6.2.

$$\omega_1 \eta_1 = -\frac{1}{3} \frac{\theta_{11}'''}{\theta_{11}'} \quad (6.13)$$

$$-\frac{1}{3} \left(\sum_{i=1}^3 e_i \right)^2 + (e_1 e_2 + e_2 e_3 + e_3 e_1) = \frac{1}{2} \left(\frac{\theta_{11}^{(5)}}{\theta_{11}^{(1)}} \right) - \frac{5}{6} \left(\frac{\theta_{11}^{(3)}}{\theta_{11}^{(1)}} \right)^2 \frac{1}{\omega_1^4}. \quad (6.14)$$

Proof. We can show the first equation (6.13) by comparing the coefficients on u^3 of (2.7).

The second equation (6.14) can be proved by comparing the coefficients on u^5 of (2.7) and by using the following equation:

$$\tilde{e}_1 \tilde{e}_2 + \tilde{e}_2 \tilde{e}_3 + \tilde{e}_3 \tilde{e}_1 = -\frac{1}{3} \left(\sum_{i=1}^3 e_i \right)^2 + (e_1 e_2 + e_2 e_3 + e_3 e_1). \quad (6.15)$$

□

Lemma 6.3. *For $\nu = 1, 2, 3$, the dependence of the period Ω on the branch points is described by the equation*

$$\frac{\partial \Omega}{\partial e_\nu} = \frac{\pi i}{\omega_1^2 \prod_{\mu \neq \nu} (e_\nu - e_\mu)}. \quad (6.16)$$

Proof. We will show the case $\nu = 1$. The other cases can be proved in the same way.

According to the definition of the Abel map, we have

$$du(x) = \frac{dx}{\sqrt{4(e - e_1)(x - e_2)(x - e_3)}}. \quad (6.17)$$

The dependence of du on e_1 is

$$\frac{\partial}{\partial e_1} \{du(x)\} = \frac{1}{2(x - e_1)} du(x). \quad (6.18)$$

Now, we consider the integral

$$\oint_{\partial \hat{E}} u(x) \frac{\partial}{\partial e_1} du = \oint_{\partial \hat{E}} \frac{1}{2(x - e_1)} u(x) du(x). \quad (6.19)$$

The left side is

$$-\omega_2 \frac{\partial \omega_1}{\partial e_1} + w_1 \frac{\partial \omega_2}{\partial e_1}. \quad (6.20)$$

We set the local coordinate of the branch point e_1 in the following way:

$$t_1 = \sqrt{x - e_1}. \quad (6.21)$$

The right side is by the residue theorem

$$\pi i \frac{du}{dt_1}(e_1) \frac{du}{dt_1}(e_1) = \frac{\pi i \omega_1^2}{\omega_1^2(e_1 - e_2)(e_1 - e_3)}. \quad (6.22)$$

By comparing the both sides, we get

$$-\frac{\omega_2}{\omega_1^2} \frac{\partial \omega_1}{\partial e_1} + \frac{\partial \omega_2}{\partial e_1} \frac{1}{\omega_1} = \frac{\pi i}{\omega_1^2(e_1 - e_2)(e_1 - e_3)}. \quad (6.23)$$

Then we obtain

$$\frac{\partial \Omega}{\partial e_1} = \frac{\pi i}{\omega_1^2(e_1 - e_2)(e_1 - e_3)}. \quad (6.24)$$

□

Lemma 6.4. For $\nu = 1, 2, 3$,

$$\frac{\partial}{\partial e_\nu} \left(\frac{\eta_1 t^2}{2\omega_1} \right) = t^2 \left(\frac{\partial}{\partial e_\nu} \log \omega_1 \right)^2 \prod_{\mu \neq \nu} (e_\nu - e_\mu) - \frac{t^2}{12}. \quad (6.25)$$

Proof. We will show the case $\nu = 1$. The other cases can be proved in the same way. From (6.13), (6.14) and the heat equation

$$\frac{\partial \theta[p, q]}{\partial z^2}(z; \Omega) = 4\pi i \frac{\partial \theta[p, q]}{\partial \Omega}(z; \Omega), \quad (6.26)$$

we get

$$\frac{\partial}{\partial e_1} \left(\frac{\eta_1 t^2}{2\omega_1} \right) = \frac{t^2}{3} \frac{\frac{\partial \omega_1}{\partial e_1}}{\omega_1^3} \frac{4\pi i \frac{\partial \theta'_{11}}{\partial \Omega}}{\theta'_{11}} - \frac{t^2}{6\omega_1^2} \frac{\partial \Omega}{\partial e_1} \frac{1}{4\pi i} \left\{ 2\omega_1^4 \left(-\frac{g_2}{4} \right) + \frac{2}{3} \left(\frac{\theta'''_{11}}{\theta'_{11}} \right)^2 \right\}.$$

By differentiating the logarithm of both sides of the formula:

$$\sqrt[4]{16(e_1 - e_2)^2(e_1 - e_3)^2(e_2 - e_3)^2} = \frac{2\pi}{\omega_1^3} (\theta'_{11})^2,$$

we get

$$3 \frac{\frac{\partial \omega_1}{\partial e_1}}{\omega_1} + \frac{1}{2} \frac{1}{(e_1 - e_2)} + \frac{1}{2} \frac{1}{(e_1 - e_3)} = 2 \frac{\frac{\partial \theta'_{11}}{\partial e_1}}{\theta'_{11}}. \quad (6.27)$$

By using (6.16) and (6.27), we obtain

$$\frac{\partial}{\partial e_1} \left(\frac{\eta_1 t^2}{2\omega_1} \right) = t^2 \left(\frac{\frac{\partial \omega_1}{\partial e_1}}{\omega_1} \right)^2 (e_1 - e_2)(e_1 - e_3) - \frac{t^2}{12}.$$

□

6.3 The tau-function

In order to compute ω_{e_ν} ($\nu = 1, 2, 3$), we will show the following lemma.

Lemma 6.5. *For $\nu = 1, 2, 3$,*

$$\begin{aligned} \text{Res}_{x=e_\nu} \frac{1}{2} \left(\frac{dY}{dx} Y^{-1}(x) \right)^2 &= -\frac{1}{8} \left(\sum_{\mu \neq \nu} \frac{1}{e_\nu - e_\mu} \right)^2 - \frac{1}{2} \frac{\partial}{\partial e_\nu} \log \omega_1 \\ &\quad + t^2 \left(\frac{\partial}{\partial e_\nu} \log \omega_1 \right)^2 \prod_{\mu \neq \nu} (e_\nu - e_\mu) + \frac{\partial}{\partial e_\nu} \left\{ \log \theta[p, q] \left(\frac{t}{\omega_1}; \Omega \right) \right\} \\ &\quad + \frac{t}{2(a - e_\nu)} \frac{\sigma[p, q]'(\frac{t}{\omega_1}; \Omega)}{\sigma[p, q](\frac{t}{\omega_1}; \Omega)} + \frac{t^2 \prod_{\mu \neq \nu} (e_\nu - e_\mu)}{4(a - e_\nu)^2} \\ &\quad + \frac{t^2}{6(a - e_\nu)} \left(\sum_{\mu \neq \nu} (e_\nu - e_\mu) \right). \end{aligned}$$

Proof. We will prove the case $\nu = 1$. The other cases can be shown in the same way. By direct calculation, we obtain

$$\frac{1}{2} \text{tr} (Y'(x) Y^{-1}(x))^2 = -\frac{\det(\Phi_x)}{\det \Phi} + \frac{1}{4} \left(\frac{(\det \Phi)_x}{\det \Phi} \right)^2.$$

We will calculate ω_1 just like ω_a in Proposition 6.1. We multiply both the numerators and the denominators of

$$\frac{\det(\Phi_x)}{\det \Phi}, \frac{(\det \Phi)_x}{\det \Phi}$$

by $\frac{1}{x_\varphi - x_\psi}$. We take the limit $P_\psi \rightarrow P_\varphi$ and get

$$\psi(P) = \frac{\partial \varphi(P)}{\partial x_\varphi}. \quad (6.28)$$

We take the limit $P_\varphi \rightarrow P$ and obtain

$$\begin{aligned}
\frac{(\det \Phi)_x}{\det \Phi} &= 2 \frac{1}{\sigma(-2u)} \lim_{P_\varphi \rightarrow P} \frac{\partial}{\partial x} \sigma(-u - u_\varphi) \\
\frac{\det(\Phi_x)}{\det \Phi} &= \frac{1}{\sigma[p, q](t)} \lim_{P_\varphi \rightarrow P} \frac{\partial^2}{\partial x \partial x_\varphi} \sigma[p, q](-u + u_\varphi + t) \\
&+ \frac{2}{\sigma[p, q](t)} \lim_{P_\varphi \rightarrow P} \frac{\partial}{\partial x} \sigma[p, q](-u + u_\varphi + t) \frac{\partial}{\partial x} \Pi(P) \\
&+ \frac{1}{\sigma(-2u)} \lim_{P_\varphi \rightarrow P} \frac{\partial^2}{\partial x \partial x_\varphi} \sigma(-u - u_\varphi) \\
&- \left(\frac{\partial}{\partial x} \Pi(P) \right)^2.
\end{aligned}$$

Therefore, we get

$$\begin{aligned}
\frac{1}{2} \operatorname{tr} (Y'(x) Y^{-1}(x))^2 &= - \lim_{P_\varphi \rightarrow P} \frac{\partial^2}{\partial x \partial x_\varphi} \log \sigma(-u - u_\varphi) \\
&- \frac{1}{\sigma[p, q](t)} \lim_{P_\varphi \rightarrow P} \frac{\partial^2}{\partial x \partial x_\varphi} \sigma[p, q](-u + u_\varphi + t) \\
&- \frac{2}{\sigma[p, q](t)} \lim_{P_\varphi \rightarrow P} \frac{\partial}{\partial x} \sigma[p, q](-u + u_\varphi + t) \frac{\partial}{\partial x} \Pi(P) \\
&+ \left(\frac{\partial}{\partial x} \Pi(P) \right)^2.
\end{aligned}$$

In order to calculate the first term of $\frac{1}{2} \operatorname{tr} (Y'(x) Y^{-1}(x))^2$, we will use the following formulas:

$$\begin{aligned}
\wp(u) &= -\frac{d^2}{du^2} \log \sigma(u) \\
\wp(u) &= x - \frac{1}{3} \sum_{\mu=1}^3 e_\mu, \quad \tilde{e}_\nu = e_\nu - \frac{1}{3} \sum_{\mu=1}^3 e_\mu \quad (\nu = 1, 2, 3) \\
\wp'(u)^2 &= 4(\wp(u) - \tilde{e}_1)(\wp(u) - \tilde{e}_2)(\wp(u) - \tilde{e}_3)
\end{aligned}$$

Then

$$\begin{aligned}
& \lim_{P_\varphi \rightarrow P} \frac{\partial^2}{\partial x \partial x_\varphi} \log \sigma(-u - u_\varphi) \\
&= \left(\frac{du}{dx} \right)^2 \frac{\sigma''(-2u)\sigma(-2u) - \sigma'(-2u)^2}{\sigma(-2u)^2} \\
&= \frac{1}{24} \sum_{\nu=1}^3 \frac{1}{(x - e_\nu)} \sum_{\mu \neq \nu} \frac{1}{(e_\nu - e_\mu)} - \frac{1}{16} \sum_{\nu=1}^3 \frac{1}{(x - e_\nu)^2}. \tag{6.29}
\end{aligned}$$

By direct calculation, we can calculate the second term of $\frac{1}{2} \text{tr}(Y'(x)Y^{-1}(x))^2$, in the following way:

$$\begin{aligned}
& \frac{1}{\sigma[p, q](t)} \lim_{P_\varphi \rightarrow P} \frac{\partial^2}{\partial x_\varphi \partial x} \sigma[p, q](-u + u_\varphi + t) \\
&= -\frac{\sigma[p, q]''(t)}{\sigma[p, q](t)} \left(\sum_{\nu=1}^3 \frac{1}{x - e_\nu} \frac{1}{4 \prod_{\mu \neq \nu} (e_\nu - e_\mu)} \right). \tag{6.30}
\end{aligned}$$

In order to calculate the third and forth terms of $\frac{1}{2} \text{tr}(Y'(x)Y^{-1}(x))^2$, we use the following relation:

$$\wp(u) = -\frac{d}{du} \zeta(u). \tag{6.31}$$

By using (6.31), we get

$$\begin{aligned}
& \frac{2}{\sigma[p, q](t)} \lim_{P_\varphi \rightarrow P} \frac{\partial}{\partial x} \sigma[p, q](-u + u_\varphi + t) \frac{\partial}{\partial x} \Pi(P) \\
&= t \frac{\sigma[p, q]'(t)}{\sigma[p, q](t)} \left\{ \frac{x}{2 \prod_{\nu=1}^3 (x - e_\nu)} + \frac{a}{2 \prod_{\nu=1}^3 (x - e_\nu)} - \frac{1}{3} \frac{\sum_{\nu=1}^3 e_\nu}{\prod_{\nu=1}^3 (x - e_\nu)} \right. \\
&\quad \left. - \frac{1}{2(x - a)^2} - \frac{1}{2} \frac{1}{(x - a)^2} \frac{\prod_{\nu=1}^3 (a - e_\nu)}{\prod_{\nu=1}^3 (x - e_\nu)} \right\} \tag{6.32}
\end{aligned}$$

and

$$\begin{aligned}
\text{Res}_{x=e_1} \left(\frac{\partial}{\partial x} \Pi(P) \right)^2 &= \frac{t^2}{36} \frac{e_1 - e_2}{e_1 - e_3} + \frac{t^2}{36} \frac{e_1 - e_3}{e_1 - e_2} + \frac{t^2}{4} \frac{(e_1 - e_2)(e_1 - e_3)}{(a - e_1)^2} \\
&\quad + \frac{t^2}{18} + \frac{t^2}{6} \frac{e_1 - e_3}{a - e_1} + \frac{t^2}{6} \frac{e_1 - e_2}{a - e_1}. \tag{6.33}
\end{aligned}$$

From (6.29), (6.30), (6.32), (6.33), we get

$$\begin{aligned}
\text{Res}_{x=e_1} \frac{1}{2} \text{tr} (Y'(x)Y^{-1}(x))^2 &= -\frac{1}{24} \sum_{j \neq 1} \frac{1}{e_1 - e_j} \\
&+ \frac{\sigma[p, q]''(t)}{\sigma[p, q](t)} \frac{1}{4(e_1 - e_2)(e_1 - e_3)} \\
&+ t \frac{\sigma[p, q]'(t)}{\sigma[p, q](t)} \left(\frac{1}{6} \sum_{j \neq 1} \frac{1}{e_1 - e_j} \right) + t \frac{\sigma[p, q]'(t)}{\sigma[p, q](t)} \frac{1}{2} \frac{1}{(a - e_1)} \\
&+ \frac{t^2}{36} \frac{e_1 - e_3}{e_1 - e_2} + \frac{t^2}{36} \frac{e_1 - e_2}{e_1 - e_3} \\
&+ \frac{t^2}{4} \frac{(e_1 - e_2)(e_1 - e_3)}{(a - e_1)^2} \\
&+ \frac{t^2}{18} + \frac{t^2}{6} \frac{e_1 - e_3}{a - e_1} + \frac{t^2}{6} \frac{e_1 - e_2}{a - e_1}.
\end{aligned}$$

By directly calculating the third term of $\text{Res}_{x=e_1} \frac{1}{2} \text{tr} (Y'(x)Y^{-1}(x))^2$, we get

$$t \frac{\sigma[p, q]'(t)}{\sigma[p, q](t)} \left(\frac{1}{6} \sum_{j \neq 1} \frac{1}{e_1 - e_j} \right) = \frac{\eta_1 t^2}{6\omega_1} \left(\frac{1}{e_1 - e_2} + \frac{1}{e_1 - e_3} \right) + \frac{t}{6\omega_1} \frac{\theta[p, q]'(\frac{t}{\omega_1}; \Omega)}{\theta[p, q](\frac{t}{\omega_1}; \Omega)} \left(\frac{1}{e_1 - e_2} + \frac{1}{e_1 - e_3} \right).$$

From Lemma 6.2, (6.26) and (6.27), we obtain

$$\begin{aligned}
\frac{\eta_1 t^2}{6\omega_1} \left(\frac{1}{e_1 - e_2} + \frac{1}{e_1 - e_3} \right) &= \frac{t^2}{6} \left(-\frac{1}{3\omega_1^2} \frac{\theta_{11}'''}{\theta_{11}'} \right) \left(\frac{1}{e_1 - e_2} + \frac{1}{e_1 - e_3} \right) \\
&= -\frac{t^2}{3} \frac{\partial \omega_1}{\partial e_1} (e_1 - e_2) - \frac{t^2}{3} \frac{\partial \omega_1}{\partial e_1} (e_1 - e_3) - \frac{t^2}{9} - \frac{t^2}{18} \frac{e_1 - e_2}{e_1 - e_3} - \frac{t^2}{18} \frac{e_1 - e_3}{e_1 - e_2}.
\end{aligned}$$

By directly calculating the second term of $\text{Res}_{x=e_1} \frac{1}{2} \text{tr} (Y'(x)Y^{-1}(x))^2$,

$$\begin{aligned}
\frac{\sigma[p, q]''(t)}{\sigma[p, q](t)} \frac{1}{4(e_1 - e_2)(e_1 - e_3)} &= \left(\frac{\eta_1}{\omega_1} \right) \frac{1}{4(e_1 - e_2)(e_1 - e_3)} + \left(\frac{\eta_1 t}{\omega_1} \right)^2 \frac{1}{4(e_1 - e_2)(e_1 - e_3)} \\
&+ \frac{2\eta_1 t}{\omega_1^2} \frac{\theta[p, q]'(\frac{t}{\omega_1}; \Omega)}{\theta[p, q](\frac{t}{\omega_1}; \Omega)} \frac{1}{4(e_1 - e_2)(e_1 - e_3)} + \frac{1}{\omega_1^2} \frac{\theta[p, q]''(\frac{t}{\omega_1}; \Omega)}{\theta[p, q](\frac{t}{\omega_1}; \Omega)} \frac{1}{4(e_1 - e_2)(e_1 - e_3)}.
\end{aligned}$$

From Lemma 6.2, (6.26) and (6.27), each of four terms of

$$\frac{\sigma[p, q]''(t)}{\sigma[p, q](t)} \frac{1}{4(e_1 - e_2)(e_1 - e_3)}$$

24

is as follows:

$$\left(\frac{\eta_1}{\omega_1}\right) \frac{1}{4(e_1 - e_2)(e_1 - e_3)} = -\frac{1}{2} \frac{\frac{\partial \omega_1}{\partial e_1}}{\omega_1} - \frac{1}{12} \frac{1}{(e_1 - e_2)} - \frac{1}{12} \frac{1}{(e_1 - e_3)}, \quad (6.34)$$

and

$$\begin{aligned} & \left(\frac{\eta_1 t}{\omega_1}\right)^2 \frac{1}{4(e_1 - e_2)(e_1 - e_3)} \\ = & t^2 \left(\frac{\frac{\partial \omega_1}{\partial e_1}}{\omega_1}\right)^2 (e_1 - e_2)(e_1 - e_3) + \frac{t^2}{36} \frac{e_1 - e_3}{e_1 - e_2} + \frac{t^2}{36} \frac{e_1 - e_2}{e_1 - e_3} \\ & + \frac{t^2}{3} \left(\frac{\frac{\partial \omega_1}{\partial e_1}}{\omega_1}\right) (e_1 - e_2) + \frac{t^2}{3} \left(\frac{\frac{\partial \omega_1}{\partial e_1}}{\omega_1}\right) (e_1 - e_3) + \frac{t^2}{18}, \end{aligned} \quad (6.35)$$

and

$$\begin{aligned} & \frac{2\eta_1 t^2}{\omega_1^2} \frac{\theta[p, q]'(\frac{t}{\omega_1}; \Omega)}{\theta[p, q](\frac{t}{\omega_1}; \Omega)} \frac{1}{4(e_1 - e_2)(e_1 - e_3)} \\ = & -\frac{\frac{\partial \omega_1}{\partial e_1}}{\omega_1^2} t \frac{\theta[p, q]'(\frac{t}{\omega_1}; \Omega)}{\theta[p, q](\frac{t}{\omega_1}; \Omega)} \\ & - \frac{t}{6\omega_1} \frac{1}{(e_1 - e_2)} \frac{\theta[p, q]'(\frac{t}{\omega_1}; \Omega)}{\theta[p, q](\frac{t}{\omega_1}; \Omega)} - \frac{t}{6\omega_1} \frac{1}{(e_1 - e_3)} \frac{\theta[p, q]'(\frac{t}{\omega_1}; \Omega)}{\theta[p, q](\frac{t}{\omega_1}; \Omega)}, \end{aligned} \quad (6.36)$$

and

$$\frac{1}{\omega_1^2} \frac{\theta[p, q]''(\frac{t}{\omega_1}; \Omega)}{\theta[p, q](\frac{t}{\omega_1}; \Omega)} \frac{1}{4(e_1 - e_2)(e_1 - e_3)} = \frac{\frac{\partial}{\partial \Omega} \theta[p, q](\frac{t}{\omega_1}; \Omega)}{\theta[p, q](\frac{t}{\omega_1}; \Omega)} \frac{\partial \Omega}{\partial e_1}. \quad (6.37)$$

We note that the sum of (6.36) and (6.37) is

$$\frac{\partial}{\partial e_1} \left\{ \log \theta[p, q] \left(\frac{t}{\omega_1}; \Omega \right) \right\} - \frac{t}{6\omega_1} \frac{1}{(e_1 - e_2)} \frac{\theta[p, q]'(\frac{t}{\omega_1}; \Omega)}{\theta[p, q](\frac{t}{\omega_1}; \Omega)} - \frac{t}{6\omega_1} \frac{1}{(e_1 - e_3)} \frac{\theta[p, q]'(\frac{t}{\omega_1}; \Omega)}{\theta[p, q](\frac{t}{\omega_1}; \Omega)}.$$

Therefore, we obtain

$$\begin{aligned}
& \text{Res}_{x=e_1} \frac{1}{2} (Y'(x)Y^{-1}(x))^2 \\
&= -\frac{1}{8} \left(\sum_{\mu \neq 1} \frac{1}{e_1 - e_\mu} \right) - \frac{1}{2} \frac{\partial}{\partial e_1} \log \omega_1 \\
&+ t^2 \left(\frac{\partial}{\partial e_1} \log \omega_1 \right)^2 \prod_{\mu \neq 1} (e_1 - e_\mu) + \frac{\partial}{\partial e_1} \left\{ \log \theta[p, q] \left(\frac{t}{\omega_1}; \Omega \right) \right\} \\
&+ \frac{t}{2(a - e_1)} \frac{\sigma[p, q]'(\frac{t}{\omega_1}; \Omega)}{\sigma[p, q](\frac{t}{\omega_1}; \Omega)} + \frac{t^2 \prod_{\mu \neq 1} (e_1 - e_\mu)}{4(a - e_1)^2} + \frac{t^2}{6(a - e_1)} \left(\sum_{\mu \neq 1} (e_1 - e_\mu) \right).
\end{aligned}$$

□

We obtain ω_a in Proposition 6.1 and get ω_{e_ν} in Lemma 6.5. Therefore, we have H_ν ($\nu = 1, 2, 3$) in the following way:

Proposition 6.6. *For $\nu = 1, 2, 3$,*

$$\begin{aligned}
H_\nu &= \frac{\partial}{\partial e_\nu} \left\{ \log \theta[p, q] \left(\frac{t}{\omega_1}; \Omega \right) \right\} - \frac{1}{2} \frac{\partial}{\partial e_\nu} \{ \log \omega_1 \} - \frac{1}{8} \sum_{\mu \neq \nu} \frac{1}{(e_\nu - e_\mu)} \\
&+ \frac{\partial}{\partial e_\nu} \left(\frac{\eta_1 t^2}{2\omega_1} \right) + \frac{\partial}{\partial e_\nu} \left\{ \frac{t^2}{4} f(e_1, e_2, e_3) \right\}.
\end{aligned}$$

Since we have calculated the Hamiltonians H_t, H_1, H_2, H_3 , we finally get the following theorem:

Theorem 6.7. *For the monodromy preserving deformation (5.1), the τ function is*

$$\tau(e_1, e_2, e_3; t) = \theta[p, q] \left(\frac{t}{\omega_1}; \Omega \right) \omega_1^{-\frac{1}{2}} \prod_{1 \leq \nu < \mu \leq 3} (e_\nu - e_\mu)^{-\frac{1}{8}} \exp\left(\frac{\eta_1 t^2}{2\omega_1}\right) \exp\left(\frac{t^2}{4} f(e_1, e_2, e_3)\right),$$

where

$$f(e_1, e_2, e_3) = -a + \frac{1}{3} \sum_{\nu=1}^3 e_\nu + \frac{1}{2} \prod_{\nu=1}^3 (a - e_\nu) \sum_{\mu=1}^3 \frac{1}{(a - e_\mu)^2}.$$

A Appendix

In [2], Miwa-Jimbo expressed the τ -function with the Weierstrass σ -function. The paper, however, is written in Japanese. We will write the summary in English. Here, we assume $e_1 + e_2 + e_3 = 0$

We fix a point α in one dimension complex torus $\mathbb{T}^1$ with period ω_1, ω_2 . We take $l \in \mathbb{Z}$ and the parameter $t \in \mathbb{C}$. We consider the following function:

$$\begin{aligned} y_l(z) &= y_l(z, t, \alpha) \\ &= \sigma(z - \alpha)^{l-1} \sigma(z + \alpha)^{-l} \sigma(z + t + (2l - 1)\alpha) \exp\left\{-\frac{t}{2} (\zeta(u - \alpha) + \zeta(u + \alpha))\right\} \end{aligned}$$

From the quasi-periodicity of the σ -function and the ζ -function, $y_l(z)$ is holomorphic in $\mathbb{T}$ except for α . Near $z = \alpha$,

$$\begin{aligned} y_l(z) &= \hat{y}_l(z) (z - \alpha)^{l-1} \exp\left(-\frac{t}{2} \left(\frac{1}{z - \alpha} - \frac{\wp''(\alpha)}{2\wp'(\alpha)}\right)\right) \\ \hat{y}_l(z) &= c_0 (1 + c_1(z - \alpha) + \cdots) \\ c_0 &= \sigma(2\alpha)^{-l} \sigma(t + 2l\alpha) \exp\left(-\frac{t}{2} \zeta(2\alpha) - \frac{t}{4} \frac{\wp''(\alpha)}{\wp'(\alpha)}\right) = c_0(t, \alpha, l) \\ c_1 &= \zeta(t + 2l\alpha) - l\zeta(2\alpha) + \frac{t}{2} \wp(2\alpha) = c_1(t, \alpha, l). \end{aligned}$$

The above property characterizes $y_l(z)$.

We set $e_1, e_2, e_3, e_\infty = \infty \in \mathbb{P}^1$ as the branch points of the covering map $\mathbb{T} \rightarrow \mathbb{P}^1, z \rightarrow x = \wp(z)$. We define the multi-valued analytic matrix $Y(x)$ in the following way:

$$\begin{aligned} Y(x) &= G_0^{-1} \tilde{Y}(z) \\ G_0^{-1} &= \text{diag}\left(\frac{1}{c_0(t, \alpha, l)}, \frac{1}{c_0(-t, \alpha, -l)}\right) \\ \tilde{Y}(z) &= \begin{pmatrix} y_l(z) & y_l(-z) \\ y_{l+1}(z) & y_{l+1}(-z) \end{pmatrix}. \end{aligned}$$

We assume $\wp(\alpha) \neq e_1, e_2, e_3, e_\infty$. Then $Y(x)$ satisfies the following linear ordinary differential equation:

$$\begin{aligned} \frac{dY}{dx} &= A(x)Y \\ A(x) &= \frac{A_{-2}}{(x - a)^2} + \frac{A_{-1}}{x - a} + \frac{B_1}{x - e_1} + \frac{B_2}{x - e_2} + \frac{B_3}{x - e_3} \\ A_{-2} &= \begin{pmatrix} \frac{\bar{t}}{2} & \\ & -\frac{\bar{t}}{2} \end{pmatrix}, \bar{t} = \wp'(\alpha)t, a = \wp(\alpha). \end{aligned}$$

In order to check this, we will study the monodromy of $Y(x)$.

A.1 The Behavior near the Irregular Singular Point

We note that

$$z - \alpha = \frac{1}{\wp'(\alpha)}(x - a) - \frac{\wp''(\alpha)}{2\wp'''(\alpha)}(x - a)^2 + \left(\frac{\wp''(\alpha)^2}{2\wp'(\alpha)^5} - \frac{\wp'''(\alpha)}{6\wp'(\alpha)^4} \right) (x - a)^3 + \cdots, \quad (\text{A.1})$$

We get the following equations:

$$\begin{aligned} Y(x) &= \hat{Y}^{(\alpha)}(x) \exp\{T^{(\alpha)}(x) + K\} \\ T^{(\alpha)}(x) &= \begin{pmatrix} \frac{\bar{t}}{2} & \\ & -\frac{\bar{t}}{2} \end{pmatrix} \frac{-1}{x - a} + \begin{pmatrix} l - 1 & \\ & -l - 1 \end{pmatrix} \log(x - a) \\ K &= \begin{pmatrix} -l + 1 & \\ & l + 1 \end{pmatrix} \log \wp'(\alpha) \\ \hat{Y}^{(\alpha)}(x) &= \left(1 + Y_1^{(\alpha)}(x - a) + \cdots \right) \\ Y_1^{(\alpha)} &= \frac{1}{\wp'(\alpha)} \tilde{Y}_1^{(\alpha)} + \frac{t}{2} \left(\frac{\wp''(\alpha)^2}{4\wp'(\alpha)^3} - \frac{\wp'''(\alpha)}{6\wp'(\alpha)^2} \right) \begin{pmatrix} 1 & \\ & -1 \end{pmatrix} - \frac{\wp''(\alpha)}{\wp'(\alpha)^2} \begin{pmatrix} l - 1 & \\ & -l - 1 \end{pmatrix} \\ \hat{Y}_1^{(\alpha)} &= \begin{pmatrix} c_1(t, \alpha, l) & \frac{c_0(-t, \alpha, 1-l)}{c_0(t, \alpha, l)} \\ \frac{c_0(t, \alpha, l+1)}{c_0(-t, \alpha, -l)} & c_1(-t, \alpha, -l) \end{pmatrix}. \end{aligned}$$

A.2 The Behavior near Regular Singular Points

We assume $\nu = 1, 2, 3$. By direct calculation, we get

$$\begin{aligned} y_l(z; t, \alpha) &= y_l(\tilde{\omega}_\nu)(1 + f_l^{(\nu)}(z - \tilde{\omega}_\nu) + \cdots) \\ y_l(\tilde{\omega}_\nu; t, \alpha) &= \frac{\sigma_\nu(t + (2l - 1)\alpha)}{\sigma_\nu(\alpha)} \\ f_l(\tilde{\omega}_\nu) &= \zeta_\nu(t + (2l - 1)\alpha) - \zeta_\nu(\tilde{\omega}_\nu + \alpha) + t\wp(\alpha + \tilde{\omega}_\nu), \end{aligned}$$

where

$$\begin{aligned} \sigma_\nu(z) &= \exp\left\{-\frac{\tilde{\eta}_\nu}{2}z\right\} \frac{\sigma(z + \tilde{\omega}_\nu)}{\sigma(\tilde{\omega}_\nu)}, \\ \zeta_\nu(z) &= \zeta(z + \tilde{\omega}_\nu) - \frac{\tilde{\eta}_\nu}{2}, \end{aligned}$$

and $\tilde{\omega}_\nu, \tilde{\eta}_\nu$ are defined in section 5. We set $G^{(\nu)}$ in the following way:

$$Y(x) = G^{(\nu)} \hat{Y}^{(\nu)}(x)(x - e_\nu) \begin{pmatrix} 0 & \\ & \frac{1}{2} \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}.$$

And

$$\begin{aligned}
G^{(\nu)} \hat{Y}^{(\nu)}(x) &= G_0^{-1} \frac{1}{2} \begin{pmatrix} y_l(z) + y_l(-z) & \frac{y_l(z) - y_l(-z)}{z - \tilde{\omega}_\nu} \\ y_{l+1}(z) + y_{l+1}(-z) & \frac{y_{l+1}(z) - y_{l+1}(-z)}{z - \tilde{\omega}_\nu} \end{pmatrix} \begin{pmatrix} 1 & \\ & \frac{z - \tilde{\omega}_\nu}{\sqrt{x - e_\nu}} \end{pmatrix} \\
&= G^{(\nu)} (1 + Y_1^{(\nu)}(x - e_\nu) + \dots) \\
G^{(\nu)} &= G_0^{-1} \begin{pmatrix} y_l(\tilde{\omega}_\nu) & \\ & y_{l+1}(\tilde{\omega}_\nu) \end{pmatrix} \begin{pmatrix} 1 & f_l^{(\nu)} \\ 1 & f_{l+1}^{(\nu)} \end{pmatrix} \begin{pmatrix} 1 & \\ & \sqrt{\frac{2}{\wp''(\tilde{\omega}_\nu)}} \end{pmatrix}
\end{aligned}$$

The case $x = \infty, z = 0$ can be calculated in the same way. By direct calculation, we get

$$\begin{aligned}
y_l(0; t, \alpha) &= (-1)^{l-1} \frac{\sigma(t + (2l-1)\alpha)}{\sigma(\alpha)} \\
f_l^{(\infty)} &= \zeta(t + (2l-1)\alpha) - (2l-1)\zeta(\alpha) + t\wp(\alpha).
\end{aligned}$$

And

$$\begin{aligned}
Y(x) &= G^{(\infty)} \hat{Y}^{(\infty)}(x) \begin{pmatrix} 1 \\ x \end{pmatrix} \begin{pmatrix} 0 & \\ & \frac{1}{2} \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \\
\hat{Y}^{(\infty)}(x) &= 1 + O\left(\frac{1}{x}\right) \\
G^{(\infty)} &= G_0^{-1} \begin{pmatrix} y_l(0) & \\ & y_{l+1}(0) \end{pmatrix} \begin{pmatrix} 1 & f_l^{(\infty)} \\ 1 & f_{l+1}^{(\infty)} \end{pmatrix}.
\end{aligned}$$

A.3 The tau-function

According to the definition of the τ -function,

$$\begin{aligned}
\frac{\partial}{\partial t} \tau(t) &= \wp'(\alpha) \operatorname{tr} Y_1^{(\alpha)} \begin{pmatrix} \frac{1}{2} & \\ & -\frac{1}{2} \end{pmatrix} \\
&= \zeta(t + 2l\alpha) + \frac{t}{2} \left(\wp(2\alpha) + \frac{\wp''(\alpha)^2}{4\wp'(\alpha)^2} - \frac{\wp'''(\alpha)}{6\wp'(\alpha)} \right) \\
&\quad - l \left(\zeta(2\alpha) + \frac{\wp''(\alpha)}{2\wp'(\alpha)} \right).
\end{aligned}$$

And

$$\begin{aligned}
\tau(t) &= \tau_l(t) = \sigma(t + 2l\alpha) \exp h_l(t) \\
h_l(t) &= \frac{t^2}{4} \left(\wp(2\alpha) + \frac{\wp''(\alpha)^2}{4\wp'(\alpha)^2} - \frac{\wp'''(\alpha)}{6\wp'(\alpha)} \right) - tl \left(\zeta(2\alpha) + \frac{\wp''(\alpha)}{2\wp'(\alpha)} \right).
\end{aligned}$$

References

- [1] L. Schlesinger, *Über eine Klasse von Differentialsystemen beliebiger Ordnung mit festen kritischen Punkten*, *J. Reine u. Angew. Math.* **141** (1912) 96-145.
- [2] M. Jimbo, T. Miwa, *Deformation of system of linear ordinary differential equations and τ -function (Japanese)*, *RIMS kokyuroku.* **388** (1980) 59-65.
- [3] M. Jimbo, T. Miwa and K. Ueno, *Monodromy preserving deformations of linear differential equations with rational coefficients*, *Physica* **1D** (1980) 80-158.
- [4] M. Jimbo and T. Miwa, *Monodromy preserving deformations of linear differential equations with rational coefficients II*, *Physica* **2D** (1981) 407-448.
- [5] D. Mumford, *Tata Lectures on Theta I*, *Progress in Math.* v.28, *Birkhauser* (1983).
- [6] A. V. Kitaev and D. A. Korotkin, *On solutions of the Schlesinger equations in terms of θ functions*, *Intern. Math. Research Notices* **17** (1998) 877-905.